

# Saving the Matter Bounce with Massive Gravity?

Based on work with **Robert Brandenberger** (McGill U.) and **Chunshan Lin** (U. of Warsaw & YITP, Kyoto U.)

JCAP **01** (2018) 011 [arXiv:1711.10472]

**Jerome Quintin**


McGill University

**Northeast Cosmology Workshop**, McGill U., Montréal

16th of March, 2018

## Introduction and Motivation

- We search for a causal mechanism that can explain the observed large scale structures of our universe from primordial fluctuations
- The standard picture is ‘horizon exit’ and ‘horizon re-entry’
- E.g., inflation:  $a(t) \propto e^{Ht}$ ,  $H \simeq \text{const.}$ ,  $H^{-1} = \text{Hubble radius}$



# Successes of Inflation

Inflation explains:


- the formation of structure problem
- the horizon problem
- the flatness problem
- the monopole problem

Also, it gives (in general):

- nearly scale-invariant power spectra of curvature and tensor perturbations
- small non-Gaussianities


# Conceptual Issues of Inflation

- Trans-Planckian problem [Brandenberger & Martin \[hep-th/0005209, hep-th/0410223\]](#)
- Singularity problem [Hawking & Penrose \[70\]; Borde & Vilenkin \[gr-qc/9612036\]; Borde \*et al.\* \[gr-qc/0110012\]](#)
- and more [Brandenberger \[1203.6698\]](#)



## Alternatives to Inflation

- There are a number of alternative scenarios for the very early universe
- E.g., **nonsingular bouncing cosmology**



# Nonsingular Bouncing Cosmology

- Solves the problems of standard Big Bang cosmology
- Free of the trans-Planckian problem
- Can avoid the initial Big Bang singularity

**What about the connection to observations?**

# Matter Bounce Scenario

- Perturbations exit the Hubble radius in a **matter-dominated contracting phase**, when  $a(\tau) \propto \tau^2$
- With a initial quantum vacuum, curvature perturbations have a scale-invariant primordial power spectrum [Wands \[gr-qc/9809062\]](#); [Finelli & Brandenberger \[hep-th/0112249\]](#)

$$v_k'' + \left( k^2 - \frac{2}{\tau^2} \right) v_k = 0, \quad ' \equiv \frac{d}{d\tau}$$

$$v_k = a \mathcal{R}_k \sqrt{2\epsilon} = \sqrt{3} a \mathcal{R}_k, \quad \epsilon \equiv -\frac{\dot{H}}{H^2} = \frac{3}{2} \left( 1 + \frac{p}{\rho} \right) = \frac{3}{2}$$

- Same for tensor modes:

$$u_k'' + \left( k^2 - \frac{2}{\tau^2} \right) u_k = 0, \quad u_k = \frac{1}{2} a h_k$$

# Power Spectra for the Matter Bounce

- Power spectra:

$$\mathcal{P}_{\mathcal{R}}(k) \equiv \frac{k^3}{2\pi^2} |\mathcal{R}_k|^2 = \frac{1}{48\pi^2} \frac{H_{B-}^2}{M_{\text{Pl}}^2}$$

$$\mathcal{P}_t(k) \equiv 2 \times \frac{k^3}{2\pi^2} |h_k|^2 = \frac{1}{2\pi^2} \frac{H_{B-}^2}{M_{\text{Pl}}^2}$$

- Tensor-to-scalar ratio:

$$r \equiv \frac{\mathcal{P}_t}{\mathcal{P}_{\mathcal{R}}} = 24$$

- Observations:  $r < 0.07$  ( $2\sigma$ ) [BICEP2 \[1510.09217\]](#)

→ **Ruled out!**



## Possible Resolution #1

- What if  $c_s \ll 1$ , e.g. with a  $k$ -essence scalar field?
- Curvature perturbations are amplified:

$$v_k'' + \left( c_s^2 k^2 - \frac{2}{\tau^2} \right) v_k = 0 \implies \mathcal{P}_{\mathcal{R}}(k) = \frac{1}{48\pi^2 c_s} \frac{H_{B-}^2}{M_{\text{Pl}}^2} \implies r = 24c_s$$

- But the scalar three-point function as well [Li, JQ et al. \[1612.02036\]](#)

$$f_{\text{NL}}^{\text{local}} \simeq -\frac{165}{16} + \frac{65}{8c_s^2}$$

- Cannot simultaneously satisfy observational bound on  $r$  and  $f_{\text{NL}}^{\text{local}} = 0.8 \pm 5.0 (1\sigma)$  [Planck \[1502.01592\]](#)
- Also,  $c_s \ll 1$  with a fluid  $\implies$  Jeans (gravitational) instability  $\implies$  black hole formation [JQ & Brandenberger \[1609.02556\]](#)

## Possible Resolution #2

- What if  $\mathcal{R}$  grows during the nonsingular bounce phase?

$$\frac{r_{\text{before bounce}}}{r_{\text{obs}}} = \left| 1 + \frac{\Delta\mathcal{R}}{\mathcal{R}_{\text{before bounce}}} \right|^2$$

- Creates large non-Gaussianities [JQ et al. \[1508.04141\]](#)

$$f_{\text{NL}} \propto \left( \frac{\Delta\mathcal{R}}{\mathcal{R}_{\text{before bounce}}} \right)^{\#}$$

- $\implies$  cannot simultaneously satisfy observational constraints on  $r$  and  $f_{\text{NL}}$

# Matter Bounce No-Go Theorem

- A lower bound on the amplification of curvature perturbations  $\mathcal{R}$ 
  - $\iff$  an upper bound on the tensor-to-scalar ratio  $r$
  - $\iff$  a lower bound on primordial non-Gaussianities  $f_{\text{NL}}$
- With Einstein gravity + a single (not necessarily canonical) scalar field:

**satisfying the current observational upper bound on  $r$  cannot be done without contradicting the current observational constraints on  $f_{\text{NL}}$  (and vice versa)** [JQ \*et al.\* \[1508.04141\]](#); [Li, JQ \*et al.\* \[1612.02036\]](#)

# Evading the No-Go Theorem

- Multiple fields, e.g., matter bounce curvaton scenario (entropy modes sourcing curvature perturbations) [Cai et al. \[1101.0822\]](#)
- Or with a single field, go beyond Einstein gravity  
→ need to modify tensor modes
- Add a nontrivial mass  $m_g$  to the graviton:

$$\mathcal{L}_{\text{tensor}}^{(2)} \supset a^2 \left[ (h'_{ij})^2 - (\nabla h_{ij})^2 \right] \rightarrow a^2 \left[ (h'_{ij})^2 - (\nabla h_{ij})^2 - m_g^2 a^2 h_{ij}^2 \right]$$
$$\implies u_k'' + \left( k^2 + m_g^2 a^2 - \frac{a''}{a} \right) u_k = 0$$

# Tensor Power Spectrum with Massive Gravity

- $m_g \ll |H_*|$  at horizon exit:

$$\mathcal{P}_t(k) \simeq \frac{1}{2\pi^2} \left( \frac{k}{aM_{\text{Pl}}} \right)^{n_t} \left| \frac{H_{B-}}{M_{\text{Pl}}} \right|^{2-n_t}, \quad n_t \simeq \frac{8m_g^2}{3H_*^2} \ll 1$$

- $m_g \gg |H_*|$  at horizon exit:

$$\mathcal{P}_t(k) \simeq \frac{2}{\pi^2} \left( \frac{k}{a} \right)^3 \frac{1}{M_{\text{Pl}}^2 m_g}$$

- In both cases,  $r(0.05 \text{ Mpc}^{-1}) \ll 0.07$  [Lin, JQ & Brandenberger \[1711.10472\]](#)

## Anisotropy problem

- Consider

$$ds^2 = -dt^2 + a^2 \sum_i e^{2\theta_i} (dx^i)^2, \quad \sum_i \theta_i = 0$$

- Einstein gravity = Friedmann equations + anisotropies:

$$\ddot{\theta}_i + 3H\dot{\theta}_i = 0 \implies \dot{\theta}_i \sim a^{-3} \implies \rho_\theta \sim \sum_i \dot{\theta}_i^2 \sim a^{-6}$$

- Analogous to a massless scalar field

$$\mathcal{L}_\theta = -\frac{1}{2} \partial_\mu \theta \partial^\mu \theta \implies p_\theta = \rho_\theta$$

- Anisotropies dominate at high energies:

$$H^2 = \frac{1}{3M_{\text{Pl}}^2} \left( \frac{\rho_{\text{m}}^0}{a^3} + \frac{\rho_{\text{rad}}^0}{a^4} \right) - \frac{k}{a^2} + \frac{\Lambda}{3} + \frac{\rho_\theta^0}{a^6}$$

# Anisotropies with Massive Gravity

- With a massive graviton [Lin, JQ & Brandenberger \[1711.10472\]](#)

$$\mathcal{L}_\theta = -\frac{1}{2}\partial_\mu\theta\partial^\mu\theta - \frac{1}{2}m_g^2\theta^2$$

$$\longrightarrow \ddot{\theta}_i + 3H\dot{\theta}_i + m_g^2\theta_i = 0, \quad \rho_\theta \sim \sum_i (\dot{\theta}_i^2 + m_g^2\theta_i^2)$$

- $m_g \gg |H| \implies \rho_\theta \sim a^{-3}$  (i.e.,  $p_\theta = 0$ )
- No anisotropy problem anymore:

$$H^2 = \frac{1}{3M_{\text{Pl}}^2} \left( \frac{\rho_m^0}{a^3} + \frac{\rho_{\text{rad}}^0}{a^4} \right) - \frac{k}{a^2} + \frac{\Lambda}{3} + \frac{\rho_\theta^0}{a^3}$$

- Would need  $m_g \gg |H_{B-}|$ , but  $m_g < 7.2 \times 10^{-23}$  eV ( $2\sigma$ ) today  
→ requires  $m_g(t)$  or symmetry breaking

# Effective Massive Gravity Action

- ADM decomposition:

$$ds^2 = -N^2 dt^2 + \gamma_{ij}(dx^i + N^i dt)(dx^j + N^j dt)$$

- Effective massive gravity action:

$$S = \int dt d^3\mathbf{x} \sqrt{\gamma} \left( N \left( \frac{M_{\text{Pl}}^2}{2} R - \Lambda \right) - m_g^2 V[\gamma_{ij}] \right)$$

- The nonderivative potential  $V[\gamma_{ij}]$  is independent of the lapse  $N$

→ **only 2 DoF propagate (2 polarization states of GWs), but diffeomorphism invariance is broken** Comelli *et al.* [1407.4991]



# Effective Massive Gravity Action

- In the EFT language, take

$$V[\gamma_{ij}] \sim M_{\text{Pl}}^2 \bar{\delta}\gamma^{ij} \bar{\delta}\gamma_{ij}$$

where  $\bar{\delta}\gamma_{ij}$  denotes the traceless part of the linear perturbations of the spatial metric [Lin & Labun \[1501.07160\]](#)

- $\implies$  background unaffected (Friedmann equations unchanged)  
scalar and vector perturbations unaffected  
tensor perturbations and anisotropies receive a mass term ( $m_g$ )
- Can be implemented with Stückelberg scalar fields (additional slides)

## Conclusions

- The matter bounce scenario is an alternative to inflation
- It naturally predicts a very large tensor-to-scalar ratio ( $r$ )
- $r$  cannot be suppressed by enhancing curvature perturbations; otherwise  $f_{\text{NL}}$  is too large
- Tensor modes can be suppressed with a (Lorentz-violating) massive graviton
- Anisotropies are suppressed likewise, but need  $m_g(t)$
- In sum, the simple idea of the matter bounce doesn't fit with observations; needs nontrivial theory to work
- Motivates us to keep looking for and testing alternative scenarios for the very early universe

# Acknowledgments

**Thank you for your attention!**

I acknowledge support from the following agencies:



Bourses d'études  
supérieures du Canada

Vanier  
Canada Graduate  
Scholarships



**NSERC**  
**CRSNG**



**McGill**

## Additional Slides: Action with Stückelberg Scalar Fields

- Introduce 1 timelike ( $\phi^0$ ) and 3 spacelike ( $\phi^i$ ) Stückelberg scalar fields with the following VEVs:

$$\phi^0 = f(t), \quad \phi^i = x^i$$

- Impose symmetries:

$$\phi^i \rightarrow \Lambda^i_j \phi^j, \quad \phi^i \rightarrow \phi^i + \Xi^i(\phi^0)$$

$\Lambda^i_j$  :  $SO(3)$  rotation operator;  $\Xi^i$  : generic function of  $\phi^0$

## Additional Slides: Action with Stückelberg Scalar Fields

- The following quantities are invariant under the symmetries [Dubovsky et al. \[hep-th/0411158\]](#)

$$X = g^{\mu\nu} \partial_\mu \phi^0 \partial_\nu \phi^0$$
$$Z^{ij} = g^{\mu\nu} \partial_\mu \phi^i \partial_\nu \phi^j - \frac{(g^{\mu\nu} \partial_\mu \phi^0 \partial_\nu \phi^i)(g^{\alpha\beta} \partial_\alpha \phi^0 \partial_\beta \phi^j)}{X}$$

- The following operator is traceless [Lin & Sasaki \[1504.01373\]](#)

$$\bar{\delta} Z^{ij} = \frac{Z^{ij}}{Z} - 3 \frac{\delta_{kl} Z^{ik} Z^{jl}}{Z^2}, \quad Z = \delta_{ij} Z^{ij}$$

- Construct quadratic operator graviton mass term:

$$\mathcal{L}_{\text{mass}} \sim M_{\text{Pl}}^2 m_g^2 \delta_{ik} \delta_{j\ell} (\bar{\delta} Z^{ij}) (\bar{\delta} Z^{k\ell})$$

- Resulting theory has 2 gravitational DoF [Lin, JQ & Brandenberger \[1711.10472\]](#)